

LEPTOSPIROSIS TESTING RECOMMENDATIONS

Jinelle Webb, DVM, MSc, DVSc, Dipl ACVIM
Mississauga-Oakville Veterinary Emergency Hospital
Jinelle.Webb@vca.com

Dog that has NOT received antibiotics and is NOT vaccinated:

- Start with Witness Lepto if you have it in clinic
- Otherwise Leptospira spp. Panel (ELISA and PCR)
- If positive on any you have your diagnosis
- If negative and clinical suspicion remains high, repeat testing or do MAT, administer appropriate antibiotics in case of false negative

Dog that has NOT received antibiotics and IS vaccinated in the previous 3 months:

- Leptospira spp. PCR
- If positive you have your diagnosis
- If negative and clinical suspicion remains high, do MAT, administer appropriate antibiotics in case of false negative on PCR

Dog that has received antibiotics and is NOT vaccinated:

- Start with Witness Lepto if you have it in clinic
- Otherwise Leptospira spp. ELISA
- If positive on any you have your diagnosis
- If negative and clinical suspicion remains high, repeat testing or do MAT

Dog that has received antibiotics and IS vaccinated in the previous 3 months:

- Acute and convalescent MAT titres

Dog that has received antibiotics and IS vaccinated but not in the previous 3 months:

- Start with Witness Lepto if you have it in clinic
- Acute and convalescent MAT titres

